

Medical Policy Manual **Draft Revision Policy: Do Not Implement**

Leuprolide Acetate Depot Suspension (Lupron Depot-Ped®)

IMPORTANT REMINDER

We develop Medical Policies to provide guidance to Members and Providers. This Medical Policy relates only to the services or supplies described in it. The existence of a Medical Policy is not an authorization, certification, explanation of benefits or a contract for the service (or supply) that is referenced in the Medical Policy. For a determination of the benefits that a Member is entitled to receive under his or her health plan, the Member's health plan must be reviewed. If there is a conflict between the medical policy and a health plan or government program (e.g., TennCare), the express terms of the health plan or government program will govern.

**The proposal is to add text/statements in red and to delete text/statements with strikethrough:
POLICY**

INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

Lupron Depot-PED is indicated for the treatment of pediatric patients with central precocious puberty (CPP).

Compendial Use

Gender dysphoria (also known as transgender and gender diverse [TGD] persons)

All other indications are considered experimental/investigational and not medically necessary.

DOCUMENTATION

Submission of the following information is necessary to initiate the prior authorization review: For central precocious puberty **initial requests**, laboratory report or medical record **documentation** of a pubertal response to a gonadotropin releasing hormone (GnRH) agonist test or a pubertal **basal** level of a ~~third-generation~~ luteinizing hormone (LH) **on a third-generation LH** assay.

PRESCRIBER SPECIALTIES

For gender dysphoria, the medication must be prescribed by or in consultation with a provider specialized in the care of transgender youth (e.g., pediatric endocrinologist, family or internal medicine physician, obstetrician-gynecologist) that has collaborated care with a mental health provider for members less than 18 years of age.

COVERAGE CRITERIA

Central Precocious Puberty (CPP)

Authorization of 12 months may be granted for treatment of CPP when all of the following criteria are met:

- **The diagnosis of CPP has been confirmed by either of the following criteria based on laboratory reference range:**
 - ~~The diagnosis of CPP has been confirmed by~~ A pubertal response to a gonadotropin releasing hormone (GnRH) agonist test ~~or a pubertal level of a third-generation luteinizing hormone (LH) assay.~~
 - **A pubertal level of luteinizing hormone (LH) on a third-generation LH assay.**
- The assessment of bone age versus chronological age supports the diagnosis of CPP.



Medical Policy Manual

Draft Revision Policy: Do Not Implement

- The member meets either of the following criteria:
 - The member is a female and was less than 8 years of age at the onset of secondary sexual characteristics.
 - The member is a male and was less than 9 years of age at the onset of secondary sexual characteristics.
- The pathologic cause of CPP has been assessed (e.g., imaging screening for intracranial tumors, genetic testing for familial CPP [e.g., MKRN3 or DLK1 **variants mutations**]).

Gender Dysphoria

*Individual is age 18 or older or the individual is less than age 18 as permissive under applicable law
Authorization of 12 months may be granted for pubertal hormonal suppression in an adolescent member when all of the following criteria are met:

- The member has a diagnosis of gender dysphoria.
- The member is able to make an informed decision to engage in treatment.
- The member has reached Tanner stage 2 of puberty or greater.
- The member's comorbid conditions are reasonably controlled.
- The member has been educated on any contraindications and side effects to therapy.
- The member has been informed of fertility preservation options.

Authorization of 12 months may be granted for gender transition when all of the following criteria are met:

- The member has a diagnosis of gender dysphoria.
- The member is able to make an informed decision to engage in treatment.
- The member will receive the requested medication concomitantly with gender-affirming hormones.
- The member's comorbid conditions are reasonably controlled.
- The member has been educated on any contraindications and side effects to therapy.
- The member has been informed of fertility preservation options.

CONTINUATION OF THERAPY

Central Precocious Puberty (CPP)

Authorization of up to 12 months may be granted for continued treatment ~~of for~~ CPP when the member meets **both** all of the following criteria:

- ~~The member is currently receiving the requested medication through a paid pharmacy or medical benefit.~~
- The member is either a female less than 12 years of age or a male less than 13 years of age.
- The member is not experiencing treatment failure (e.g., clinical pubertal progression, lack of growth deceleration, continued excessive bone age advancement).

Gender Dysphoria

*Individual is age 18 or older or the individual is less than age 18 as permissive under applicable law
Authorization of 12 months may be granted for continued treatment for pubertal hormonal suppression in adolescent members requesting reauthorization when all of the following criteria are met:

- The member has a diagnosis of gender dysphoria.
- The member is able to make an informed decision to engage in treatment.
- The member has previously reached Tanner stage 2 of puberty or greater.
- The member's comorbid conditions are reasonably controlled.

Medical Policy Manual **Draft Revision Policy: Do Not Implement**

- The member has been educated on any contraindications and side effects to therapy.
- Before the start of therapy, the member has been informed of fertility preservation options.

Authorization of 12 months may be granted for continued treatment for gender transition in members requesting reauthorization when all of the following criteria are met:

- The member has a diagnosis of gender dysphoria.
- The member is able to make an informed decision to engage in treatment.
- The member will receive the requested medication concomitantly with gender-affirming hormones.
- The member's comorbid conditions are reasonably controlled.
- The member has been educated on any contraindications and side effects to therapy.
- Before the start of therapy, the member has been informed of fertility preservation options.

OTHER

Per state regulatory guidelines around gender dysphoria, age restrictions may apply.

APPLICABLE TENNESSEE STATE MANDATE REQUIREMENTS

BlueCross BlueShield of Tennessee's Medical Policy complies with Tennessee Code Annotated Section 56-7-2352 regarding coverage of off-label indications of Food and Drug Administration (FDA) approved drugs when the off-label use is recognized in one of the statutorily recognized standard reference compendia or in the published peer-reviewed medical literature.

ADDITIONAL INFORMATION

For appropriate chemotherapy regimens, dosage information, contraindications, precautions, warnings, and monitoring information, please refer to one of the standard reference compendia (e.g., the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) published by the National Comprehensive Cancer Network®, Drugdex Evaluations of Micromedex Solutions at Truven Health, or The American Hospital Formulary Service Drug Information).

REFERENCES

1. Lupron Depot-PED [package insert]. North Chicago, IL: AbbVie Inc.; **November 2025**.
2. Kletter GB, Klein KO, Wong YY. A pediatrician's guide to central precocious puberty. *Clin Pediatr*. 2015;54:414-424.
3. Carel JC, Eugster EA, Rogol A, et al. Consensus statement on the use of gonadotropin-releasing hormone analogs in children. *Pediatrics*. 2009;123:e752-e762.
4. Bangalore Krishna K, Fuqua JS, Rogol AD, et al. Use of gonadotropin-releasing hormone analogs in children: Update by an international consortium. *Horm Res Paediatr*. 2019;91(6):357-372.
5. Bangalore Krishna K, Silverman LA. Diagnosis of central precocious puberty. *Endocrinol Metab Clin North Am*. 2024;53(2):217-227.
6. Kaplowitz P, Bloch C ~~the Section on Endocrinology~~. Evaluation and referral of children with signs of early puberty. *Pediatrics*. 2016;137(1). doi:10.1542/peds.2015-3732 ~~:e20153732~~.
7. **Cheuiche AV, da Silveira LG, de Paula LCP, et al. Diagnosis and management of precocious sexual maturation: an updated review. *Eur J Pediatr*. 2021;180(10):3073-3087.**
8. **Popovic J, Geffner ME, Rogol AD, et al. Gonadotropin-releasing hormone analog therapies for children with central precocious puberty in the United States. *Front Pediatr*. 2022;10:968485. doi:10.3389/fped.2022.968485**



9. Hembree WC, Cohen-Kettenis PT, Gooren L, et al. Endocrine Treatment of Gender-Dysphoric/Gender-Incongruent Persons: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab.* 2017;102(11):3869–3903.
10. Gender Identity Research and Education Society. Guidance for GPs and other clinicians on the treatment of gender variant people. UK Department of Health. Published March 10, 2008.
11. Coleman E, Radix AE, Brown GR, et al. Standards of care for the health of transgender, and gender diverse people, version 8. 2022 ;23 (Suppl 1): S1-S259. doi: 10.1080/26895269.2022.2100644
12. ~~Cheuiche AV, da Silveira LG, de Paula LCP, et al. Diagnosis and management of precocious sexual maturation: an updated review. *Eur J Pediatr.* 2021;180(10):3073–3087.~~
13. Mahfouda S, Moore JK, Siafarikas A, et al. Puberty Suppression in transgender children and adolescents. *Lancet Diabetes Endocrinol.* 2017; 5: 816-26.
14. Health Care for Transgender and Gender Diverse Individuals. ©2021 The American College of Obstetricians and Gynecologists. Available at: <https://www.acog.org/clinical/clinical-guidance/committee-opinion/articles/2021/03/health-care-for-transgender-and-gender-diverse-individuals>.
15. ~~Popovic J, Geffner ME, Rogol AD, et al. Gonadotropin-releasing hormone analog therapies for children with central precocious puberty in the United States. *Front Pediatr.* 2022;10:968485. doi:10.3389/fped.2022.968485~~

EFFECTIVE DATE

ID_CHS_2026