

## Medical Policy Manual

## Draft Revision Policy: Do Not Implement

**Pegfilgrastim (Neulasta®); Pegfilgrastim-unne ( Armlupeg® ); Pegfilgrastim-pccg (Ennumo™); Pegfilgrastim-jmdb (Fulphila®); Pegfilgrastim-pbbk (Fynetra®); Pegfilgrastim-apgf (Nyvepria™); Pegfilgrastim-fpgk (Stimufend®); Pegfilgrastim-cbqv (Udenyca®); Pegfilgrastim-bmez (Ziextenzo™)**

Some agents on this policy may require step therapy See “Step Therapy Requirements for Provider Administered Specialty Medications” Document at:

[https://www.bcbst.com/docs/providers/Comm\\_BC\\_PAD\\_Step\\_Therapy\\_Guide.pdf](https://www.bcbst.com/docs/providers/Comm_BC_PAD_Step_Therapy_Guide.pdf)

### IMPORTANT REMINDER

We develop Medical Policies to provide guidance to Members and Providers. This Medical Policy relates only to the services or supplies described in it. The existence of a Medical Policy is not an authorization, certification, explanation of benefits or a contract for the service (or supply) that is referenced in the Medical Policy. For a determination of the benefits that a Member is entitled to receive under his or her health plan, the Member's health plan must be reviewed. If there is a conflict between the medical policy and a health plan or government program (e.g., TennCare), the express terms of the health plan or government program will govern.

**The proposal is to add text/statements in red and to delete text/statements with strikethrough:**

### POLICY

### INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

#### FDA-Approved Indications

##### **Neulasta**

##### **Patients with Cancer Receiving Myelosuppressive Chemotherapy**

Neulasta is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.

##### **Hematopoietic Subsyndrome of Acute Radiation Syndrome**

Neulasta is indicated to increase survival in patients acutely exposed to myelosuppressive doses of radiation (Hematopoietic Subsyndrome of Acute Radiation Syndrome).

##### **Armlupeg**

##### **Patients with Cancer Receiving Myelosuppressive Chemotherapy**

Armlupeg is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.

##### **Hematopoietic Subsyndrome of Acute Radiation Syndrome**

Armlupeg is indicated to increase survival in patients acutely exposed to myelosuppressive doses of radiation (Hematopoietic Subsyndrome of Acute Radiation Syndrome).

##### **Ennumo**

## Medical Policy Manual

## **Draft Revision Policy: Do Not Implement**

### **Patients with Cancer Receiving Myelosuppressive Chemotherapy**

Ennumo is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.

### **Hematopoietic Subsyndrome of Acute Radiation Syndrome**

Ennumo is indicated to increase survival in patients acutely exposed to myelosuppressive doses of radiation (Hematopoietic Subsyndrome of Acute Radiation Syndrome).

### **Fulphila**

#### **Patients with Cancer Receiving Myelosuppressive Chemotherapy**

Fulphila is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia

### **Udenyca**

#### **Patients with Cancer Receiving Myelosuppressive Chemotherapy**

Udenyca is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.

### **Hematopoietic Subsyndrome of Acute Radiation Syndrome**

Udenyca is indicated to increase survival in patients acutely exposed to myelosuppressive doses of radiation.

### **Ziextenzo**

#### **Patients with Cancer Receiving Myelosuppressive Chemotherapy**

Ziextenzo is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.

### **Hematopoietic Subsyndrome of Acute Radiation Syndrome**

Ziextenzo is indicated to increase survival in patients acutely exposed to myelosuppressive doses of radiation.

### **Nyvepria**

#### **Patients with Cancer Receiving Myelosuppressive Chemotherapy**

Nyvepria is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.

### **Fylnetra**

#### **Patients with Cancer Receiving Myelosuppressive Chemotherapy**

Fylnetra is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.

### **Hematopoietic Subsyndrome of Acute Radiation Syndrome**

Fylnetra is indicated to increase survival in patients acutely exposed to myelosuppressive doses of radiation.

### **Stimufend**

#### **Patients with Cancer Receiving Myelosuppressive Chemotherapy**

## Medical Policy Manual

## **Draft Revision Policy: Do Not Implement**

Stimufend is indicated to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive anti-cancer drugs associated with a clinically significant incidence of febrile neutropenia.

### **Hematopoietic Subsyndrome of Acute Radiation Syndrome**

Stimufend is indicated to increase survival in patients acutely exposed to myelosuppressive doses of radiation.

#### Compendial Use

- Stem cell transplantation-related indications
- Prophylaxis for chemotherapy-induced febrile neutropenia in patients with solid tumors
- Hematopoietic Acute Radiation Syndrome
- Hairy cell leukemia, neutropenic fever

All other indications are considered experimental/investigational and not medically necessary.

## **DOCUMENTATION**

### **Primary Prophylaxis of Febrile Neutropenia**

- Documentation must be provided of the member's diagnosis and chemotherapeutic regimen.
- If chemotherapeutic regimen has a low or intermediate risk of febrile neutropenia (20% and less), documentation must be provided outlining the member's risk factors that confirm the member is at high risk for febrile neutropenia.

## **COVERAGE CRITERIA**

### **Prevention of Neutropenia in Cancer Patients Receiving Myelosuppressive Chemotherapy**

Authorization of 6 months may be granted for prevention of febrile neutropenia when all of the following criteria are met:

- The requested medication will not be used in combination with other colony stimulating factors within any chemotherapy cycle.
- The member will not receive chemotherapy at the same time as they receive radiation therapy.
- The requested medication will not be administered with weekly chemotherapy regimens.
- One of the following criteria is met:
  - The requested medication will be used for primary prophylaxis in members with a solid tumor or non-myeloid malignancies who have received, are currently receiving, or will be receiving any of the following:
    - Myelosuppressive anti-cancer therapy that is expected to result in greater than 20% incidence of febrile neutropenia (FN) (See Appendix A).
    - Myelosuppressive anti-cancer therapy that is expected to result in 10 – 20% risk of FN (See Appendix B) and who are considered to be at high risk of FN because of bone marrow compromise, co-morbidities, or other patient specific risk factors (See Appendix C).
    - Myelosuppressive anti-cancer therapy that is expected to result in less than 10% risk of FN and who have at least 2 patient-related risk factors (See Appendix C).
  - The requested medication will be used for secondary prophylaxis in members with solid tumors or non-myeloid malignancies who experienced a febrile neutropenic complication or a dose-limiting

## Medical Policy Manual

## **Draft Revision Policy: Do Not Implement**

neutropenic event (a nadir or day of treatment count impacting the planned dose of chemotherapy) from a prior cycle of similar chemotherapy, with the same dose and scheduled planned for the current cycle (for which primary prophylaxis was not received).

### **OTHER INDICATIONS**

Authorization of 6 months may be granted for members with any of the following indications:

- Stem cell transplantation-related indications
  - Hematopoietic Subsyndrome of Acute Radiation Syndrome
  - Treatment for radiation-induced myelosuppression following a radiological/nuclear incident
  - Hairy cell leukemia
- Members with hairy cell leukemia with neutropenic fever following chemotherapy

### **CONTINUATION OF THERAPY**

All members (including new members) requesting authorization for continuation of therapy must meet all requirements in the coverage criteria.

### **APPENDIX**

#### **APPENDIX A: Selected Chemotherapy Regimens with an Incidence of Febrile Neutropenia of Greater than 20%**

This list is not comprehensive; there are other agents/regimens that have an intermediate/high risk for development of febrile neutropenia.

##### **Acute Lymphoblastic Leukemia**

Select ALL regimens as directed by treatment protocol (see NCCN guidelines ALL)

##### **Bladder Cancer**

Dose dense MVAC (methotrexate, vinblastine, doxorubicin, cisplatin)

##### **Bone Cancer**

- VAIA (vincristine, doxorubicin, ifosfamide, and dactinomycin)
- VDC-IE (vincristine, doxorubicin or dactinomycin, and cyclophosphamide alternating with ifosfamide and etoposide)
- Cisplatin/doxorubicin
- VDC (cyclophosphamide, vincristine, doxorubicin or dactinomycin)
- VIDE (vincristine, ifosfamide, doxorubicin or dactinomycin, etoposide)

##### **Breast Cancer**

- Dose-dense AC (doxorubicin, cyclophosphamide) followed by dose-dense paclitaxel
- TAC (docetaxel, doxorubicin, cyclophosphamide)
- TC (docetaxel, cyclophosphamide)
- TCH (docetaxel, carboplatin, trastuzumab)

##### **Head and Neck Squamous Cell Carcinoma**

TPF (docetaxel, cisplatin, 5-fluorouracil)

##### **Hodgkin Lymphoma**

## Medical Policy Manual

## **Draft Revision Policy: Do Not Implement**

- Brentuximab vedotin + AVD (doxorubicin, vinblastine, dacarbazine)
- Nivolumab + AVD (doxorubicin, vinblastine, dacarbazine)
- Escalated BEACOPP (bleomycin, etoposide, doxorubicin, cyclophosphamide, vincristine, procarbazine, prednisone)
- BrECADD (brentuximab vedotin, etoposide, cyclophosphamide, doxorubicin, dacarbazine, dexamethasone)

### **Kidney Cancer**

Doxorubicin/gemcitabine

### **Non-Hodgkin's Lymphoma**

- CHP (cyclophosphamide, doxorubicin, prednisone) + brentuximab vedotin
- Dose-adjusted EPOCH (etoposide, prednisone, vincristine, cyclophosphamide, doxorubicin) ± rituximab
- ICE (ifosfamide, carboplatin, etoposide) ± rituximab
- Dose-dense CHOP-14 (cyclophosphamide, doxorubicin, vincristine, prednisone) ± rituximab
- CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone)
- MINE (mesna, ifosfamide, mitoxantrone, etoposide) ± rituximab
- DHAP (dexamethasone, cisplatin, cytarabine) ± rituximab
- ESHAP (etoposide, methylprednisolone, cisplatin, cytarabine) ± rituximab
- HyperCVAD ± rituximab (cyclophosphamide, vincristine, doxorubicin, dexamethasone ± rituximab)
- Pola-R-CHP (polatuzumab vedotin-piiq, rituximab, cyclophosphamide, doxorubicin, prednisone)

### **Melanoma**

Dacarbazine-based combination with IL-2, interferon alpha (dacarbazine, cisplatin, vinblastine, IL-2, interferon alfa)

### **Multiple Myeloma**

- VTD-PACE (dexamethasone/thalidomide/cisplatin/doxorubicin/cyclophosphamide/etoposide + bortezomib)
- DT-PACE (dexamethasone/thalidomide/cisplatin/doxorubicin/cyclophosphamide/etoposide)

### **Ovarian Cancer**

- Topotecan ± bevacizumab
- Docetaxel
- Carboplatin/docetaxel

### **Soft Tissue Sarcoma**

- MAID (mesna, doxorubicin, ifosfamide, dacarbazine)
- Doxorubicin
- Ifosfamide/doxorubicin

### **Small Cell Lung Cancer**

Topotecan

### **Testicular Cancer**

- VeIP (vinblastine, ifosfamide, cisplatin)
- VIP (etoposide, ifosfamide, cisplatin)
- TIP (paclitaxel, ifosfamide, cisplatin)

### **Gestational Trophoblastic Neoplasia**

- EMA/EP (etoposide, methotrexate, dactinomycin/etoposide, cisplatin)

## Medical Policy Manual

## **Draft Revision Policy: Do Not Implement**

- EP/EMA (etoposide, cisplatin/etoposide, methotrexate, dactinomycin)
- TP/TE (paclitaxel, cisplatin/paclitaxel, etoposide)
- BEP (bleomycin, etoposide, cisplatin)
- TIP (Paclitaxel, ifosfamide, cisplatin)
- VIP (etoposide, ifosfamide, cisplatin)
- ICE (ifosfamide, carboplatin, etoposide)

### **Wilms Tumor**

- Regimen M (vincristine, dactinomycin, doxorubicin, cyclophosphamide, etoposide)
- Regimen I (vincristine, doxorubicin, cyclophosphamide, etoposide)
- Revised Regimen UH-1 (vincristine, doxorubicin, cyclophosphamide, carboplatin, etoposide)
- Revised Regimen UH-2 (vincristine, doxorubicin, cyclophosphamide, carboplatin, etoposide, irinotecan)

Applies to chemotherapy regimens with or without monoclonal antibodies (e.g., trastuzumab, rituximab)

### **APPENDIX B: Selected Chemotherapy Regimens with an Incidence of Febrile Neutropenia of 10% to 20%**

This list is not comprehensive; there are other agents/regimens that have an intermediate/high risk for development of febrile neutropenia.

### **Occult Primary – Adenocarcinoma**

Gemcitabine/docetaxel

### **Breast Cancer**

- Docetaxel ± trastuzumab
- AC (doxorubicin, cyclophosphamide) + sequential docetaxel (taxane portion only)
- AC + sequential docetaxel + trastuzumab
- Paclitaxel every 21 days ± trastuzumab
- Sacituzumab govitecan-hziy
- TC (docetaxel, cyclophosphamide)

### **Cervical Cancer**

- Irinotecan
- Cisplatin/topotecan
- Paclitaxel/cisplatin ± bevacizumab
- Topotecan

### **Colorectal Cancer**

FOLFIRINOX (fluorouracil, leucovorin, oxaliplatin, irinotecan)

### **Esophageal and Gastric Cancers**

Irinotecan/cisplatin

### **Non-Hodgkin's Lymphomas**

- GDP (gemcitabine, dexamethasone, cisplatin/carboplatin)
- GDP (gemcitabine, dexamethasone, cisplatin/carboplatin) + rituximab
- CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) including regimens with pegylated liposomal doxorubicin
- CHOP + rituximab (cyclophosphamide, doxorubicin, vincristine, prednisone, rituximab) including regimens with pegylated liposomal doxorubicin

## Medical Policy Manual

**Draft Revision Policy: Do Not Implement**

- Bendamustine

### **Non-Small Cell Lung Cancer**

- Cisplatin/paclitaxel
- Cisplatin/vinorelbine
- Cisplatin/docetaxel
- Cisplatin/etoposide
- Carboplatin/paclitaxel
- Docetaxel

### **Pancreatic Cancer**

FOLFIRINOX (fluorouracil, leucovorin, oxaliplatin, irinotecan)

### **Prostate Cancer**

Cabazitaxel

### **Small Cell Lung Cancer**

Etoposide/carboplatin

### **Testicular Cancer**

- BEP (bleomycin, etoposide, cisplatin)
- Etoposide/cisplatin

### **Uterine Sarcoma**

Docetaxel

Applies to chemotherapy regimens with or without monoclonal antibodies (e.g., trastuzumab, rituximab)

## **APPENDIX C: Patient Risk Factors**

This list is not all-inclusive.

- Active infections, open wounds, or recent surgery
- Age greater than or equal to 65 years
- Bone marrow involvement by tumor producing cytopenias
- Previous chemotherapy or radiation therapy
- Poor nutritional status
- Poor performance status
- Previous episodes of FN
- Other serious co-morbidities, including renal dysfunction, liver dysfunction, HIV infection, cardiovascular disease
- Persistent neutropenia

## **MEDICATION QUANTITY LIMITS**

| <b>Drug Name</b>                 | <b>Diagnosis</b>                                         | <b>Maximum Dosing Regimen</b>                                                                         |
|----------------------------------|----------------------------------------------------------|-------------------------------------------------------------------------------------------------------|
| Armlupeg<br>(Pegfilgrastim-unne) | Hematopoietic Subsyndrome of<br>Acute Radiation Syndrome | Route of Administration: Subcutaneous<br>≥18 year(s)<br>6mg every week for 2 doses<br><br>≤18 year(s) |



|                                  |                                                       |                                                                                                                                                                                                                                                                                                                                                                  |
|----------------------------------|-------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                  |                                                       | <p>≥45kg<br/>6mg every week for 2 doses</p>                                                                                                                                                                                                                                                                                                                      |
| Armlupeg<br>(Pegfilgrastim-unne) | Prevention of Febrile Neutropenia                     | <p>Route of Administration: Subcutaneous<br/>≥18 year(s)<br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>&lt;18 year(s)<br/>≥45kg<br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> |
| Armlupeg<br>(Pegfilgrastim-unne) | Stem Cell Transplantation-Related Indications         | <p>Route of Administration: Subcutaneous<br/>6mg once</p>                                                                                                                                                                                                                                                                                                        |
| Ennumo<br>(Pegfilgrastim-pccg)   | Hematopoietic Subsyndrome of Acute Radiation Syndrome | <p>Route of Administration: Subcutaneous<br/>≥18 year(s)<br/>6mg every week for 2 doses</p> <p>&lt;18 year(s)<br/>≥45kg<br/>6mg every week for 2 doses</p>                                                                                                                                                                                                       |
| Ennumo<br>(Pegfilgrastim-pccg)   | Prevention of Febrile Neutropenia                     | <p>Route of Administration: Subcutaneous<br/>≥18 year(s)<br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>&lt;18 year(s)<br/>≥45kg<br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> |
| Ennumo<br>(Pegfilgrastim-pccg)   | Stem Cell Transplantation-Related Indications         | <p>Route of Administration: Subcutaneous<br/>6mg once</p>                                                                                                                                                                                                                                                                                                        |
| Fulphila<br>(Pegfilgrastim-jmdb) | Hematopoietic Subsyndrome of Acute Radiation Syndrome | <p>Route of Administration: Subcutaneous<br/>≥18 year(s)<br/>6mg every week for 2 doses</p> <p>&lt;18 year(s)<br/>≥45kg<br/>6mg every week for 2 doses</p> <p>31 - &lt;45kg<br/>4mg every week for 2 doses</p> <p>21 - &lt;31kg<br/>2.5mg every week for 2 doses</p> <p>10 - &lt;21kg</p>                                                                        |



|                                  |                                                       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|----------------------------------|-------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                  |                                                       | <p>1.5mg every week for 2 doses</p> <p>&lt;10kg<br/>0.1mg/kg every week for 2 doses</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Fulphila<br>(Pegfilgrastim-jmdb) | Prevention of Febrile Neutropenia                     | <p>Route of Administration: Subcutaneous</p> <p><u>≥18 year(s)</u><br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p><u>&lt;18 year(s)</u><br/>≥45kg<br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>31 - &lt;45kg<br/>4mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>21 - &lt;31kg<br/>2.5mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>10 - &lt;21kg<br/>1.5mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>&lt;10kg<br/>0.1mg/kg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> |
| Fulphila<br>(Pegfilgrastim-jmdb) | Stem Cell Transplantation-Related Indications         | <p>Route of Administration: Subcutaneous</p> <p>6mg once</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Fylnetra<br>(Pegfilgrastim-pbbk) | Hematopoietic Subsyndrome of Acute Radiation Syndrome | <p>Route of Administration: Subcutaneous</p> <p><u>≥18 year(s)</u><br/>6mg every week for 2 doses</p> <p><u>&lt;18 year(s)</u><br/>≥45kg<br/>6mg every week for 2 doses</p> <p>31 - &lt;45kg<br/>4mg every week for 2 doses</p> <p>21 - &lt;31kg<br/>2.5mg every week for 2 doses</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |



|                                  |                                                       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|----------------------------------|-------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                  |                                                       | <p>10 - &lt;21kg<br/>1.5mg every week for 2 doses</p> <p>&lt;10kg<br/>0.1mg/kg every week for 2 doses</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Fylnetra<br>(Pegfilgrastim-pbbk) | Prevention of Febrile Neutropenia                     | <p>Route of Administration: Subcutaneous<br/><u>≥18 year(s)</u><br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p><u>&lt;18 year(s)</u><br/>≥45kg<br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>31 - &lt;45kg<br/>4mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>21 - &lt;31kg<br/>2.5mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>10 - &lt;21kg<br/>1.5mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>&lt;10kg<br/>0.1mg/kg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> |
| Fylnetra<br>(Pegfilgrastim-pbbk) | Stem Cell Transplantation-Related Indications         | <p>Route of Administration: Subcutaneous<br/>6mg once</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Neulasta<br>(Pegfilgrastim)      | Hematopoietic Subsyndrome of Acute Radiation Syndrome | <p>Route of Administration: Subcutaneous<br/><u>≥18 year(s)</u><br/>6mg every week for 2 doses</p> <p><u>&lt;18 year(s)</u><br/>≥45kg<br/>6mg every week for 2 doses</p> <p>31 - &lt;45kg<br/>4mg every week for 2 doses</p> <p>21 - &lt;31kg<br/>2.5mg every week for 2 doses</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |



|                                  |                                                       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|----------------------------------|-------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                  |                                                       | <p>10 - &lt;21kg<br/>1.5mg every week for 2 doses</p> <p>&lt;10kg<br/>0.1mg/kg every week for 2 doses</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Neulasta<br>(Pegfilgrastim)      | Prevention of Febrile Neutropenia                     | <p>Route of Administration: Subcutaneous</p> <p><u>≥18 year(s)</u><br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p><u>&lt;18 year(s)</u><br/>≥45kg<br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>31 - &lt;45kg<br/>4mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>21 - &lt;31kg<br/>2.5mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>10 - &lt;21kg<br/>1.5mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>&lt;10kg<br/>0.1mg/kg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> |
| Neulasta<br>(Pegfilgrastim)      | Stem Cell Transplantation-Related Indications         | <p>Route of Administration: Subcutaneous</p> <p>6mg once</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Nyvepria<br>(Pegfilgrastim-apgf) | Hematopoietic Subsyndrome of Acute Radiation Syndrome | <p>Route of Administration: Subcutaneous</p> <p><u>≥18 year(s)</u><br/>6mg every week for 2 doses</p> <p><u>&lt;18 year(s)</u><br/>≥45kg<br/>6mg every week for 2 doses</p> <p>31 - &lt;45kg<br/>4mg every week for 2 doses</p> <p>21 - &lt;31kg<br/>2.5mg every week for 2 doses</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |



|                                   |                                                       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-----------------------------------|-------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                   |                                                       | <p>10 - &lt;21kg<br/>1.5mg every week for 2 doses</p> <p>&lt;10kg<br/>0.1mg/kg every week for 2 doses</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Nyvepria<br>(Pegfilgrastim-apgf)  | Prevention of Febrile Neutropenia                     | <p>Route of Administration: Subcutaneous</p> <p><u>≥18 year(s)</u><br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p><u>&lt;18 year(s)</u><br/>≥45kg<br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>31 - &lt;45kg<br/>4mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>21 - &lt;31kg<br/>2.5mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>10 - &lt;21kg<br/>1.5mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>&lt;10kg<br/>0.1mg/kg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> |
| Nyvepria<br>(Pegfilgrastim-apgf)  | Stem Cell Transplantation-Related Indications         | <p>Route of Administration: Subcutaneous</p> <p>6mg once</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Stimufend<br>(Pegfilgrastim-fpgk) | Hematopoietic Subsyndrome of Acute Radiation Syndrome | <p>Route of Administration: Subcutaneous</p> <p><u>≥18 year(s)</u><br/>6mg every week for 2 doses</p> <p><u>&lt;18 year(s)</u><br/>≥45kg<br/>6mg every week for 2 doses</p> <p>31 - &lt;45kg<br/>4mg every week for 2 doses</p> <p>21 - &lt;31kg<br/>2.5mg every week for 2 doses</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |



|                                   |                                                       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-----------------------------------|-------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                   |                                                       | <p>10 - &lt;21kg<br/>1.5mg every week for 2 doses</p> <p>&lt;10kg<br/>0.1mg/kg every week for 2 doses</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Stimufend<br>(Pegfilgrastim-fpgk) | Prevention of Febrile Neutropenia                     | <p>Route of Administration: Subcutaneous</p> <p><u>≥18 year(s)</u><br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p><u>&lt;18 year(s)</u><br/>≥45kg<br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>31 - &lt;45kg<br/>4mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>21 - &lt;31kg<br/>2.5mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>10 - &lt;21kg<br/>1.5mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>&lt;10kg<br/>0.1mg/kg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> |
| Stimufend<br>(Pegfilgrastim-fpgk) | Stem Cell Transplantation-Related Indications         | <p>Route of Administration: Subcutaneous</p> <p>6mg once</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Udenyca<br>(Pegfilgrastim-cbqv)   | Hematopoietic Subsyndrome of Acute Radiation Syndrome | <p>Route of Administration: Subcutaneous</p> <p><u>≥18 year(s)</u><br/>6mg every week for 2 doses</p> <p><u>&lt;18 year(s)</u><br/>≥45kg<br/>6mg every week for 2 doses</p> <p>31 - &lt;45kg<br/>4mg every week for 2 doses</p> <p>21 - &lt;31kg<br/>2.5mg every week for 2 doses</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |



|                                   |                                                       |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-----------------------------------|-------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                   |                                                       | <p>10 - &lt;21kg<br/>1.5mg every week for 2 doses</p> <p>&lt;10kg<br/>0.1mg/kg every week for 2 doses</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Udenyca<br>(Pegfilgrastim-cbqv)   | Prevention of Febrile Neutropenia                     | <p>Route of Administration: Subcutaneous</p> <p><u>≥18 year(s)</u><br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p><u>&lt;18 year(s)</u><br/>≥45kg<br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>31 - &lt;45kg<br/>4mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>21 - &lt;31kg<br/>2.5mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>10 - &lt;21kg<br/>1.5mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>&lt;10kg<br/>0.1mg/kg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> |
| Udenyca<br>(Pegfilgrastim-cbqv)   | Stem Cell Transplantation-Related Indications         | <p>Route of Administration: Subcutaneous</p> <p>6mg once</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Ziextenzo<br>(Pegfilgrastim-bmez) | Hematopoietic Subsyndrome of Acute Radiation Syndrome | <p>Route of Administration: Subcutaneous</p> <p><u>≥18 year(s)</u><br/>6mg every week for 2 doses</p> <p><u>&lt;18 year(s)</u><br/>≥45kg<br/>6mg every week for 2 doses</p> <p>31 - &lt;45kg<br/>4mg every week for 2 doses</p> <p>21 - &lt;31kg<br/>2.5mg every week for 2 doses</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |



|                                   |                                               |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|-----------------------------------|-----------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                   |                                               | <p>10 - &lt;21kg<br/>1.5mg every week for 2 doses</p> <p>&lt;10kg<br/>0.1mg/kg every week for 2 doses</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Ziextenzo<br>(Pegfilgrastim-bmez) | Prevention of Febrile Neutropenia             | <p>Route of Administration: Subcutaneous</p> <p><u>≥18 year(s)</u><br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p><u>&lt;18 year(s)</u><br/>≥45kg<br/>6mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>31 - &lt;45kg<br/>4mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>21 - &lt;31kg<br/>2.5mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>10 - &lt;21kg<br/>1.5mg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> <p>&lt;10kg<br/>0.1mg/kg once per chemotherapy cycle; do not administer between 14 days prior to and 24 hours after administration of chemotherapy</p> |
| Ziextenzo<br>(Pegfilgrastim-bmez) | Stem Cell Transplantation-Related Indications | <p>Route of Administration: Subcutaneous</p> <p>6mg once</p>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |

## APPLICABLE TENNESSEE STATE MANDATE REQUIREMENTS

BlueCross BlueShield of Tennessee's Medical Policy complies with Tennessee Code Annotated Section 56-7-2352 regarding coverage of off-label indications of Food and Drug Administration (FDA) approved drugs when the off-label use is recognized in one of the statutorily recognized standard reference compendia or in the published peer-reviewed medical literature.

## ADDITIONAL INFORMATION

For appropriate chemotherapy regimens, dosage information, contraindications, precautions, warnings, and monitoring information, please refer to one of the standard reference compendia (e.g., the NCCN Clinical Practice

## Medical Policy Manual

## **Draft Revision Policy: Do Not Implement**

Guidelines in Oncology (NCCN Guidelines®) published by the National Comprehensive Cancer Network®, Drugdex Evaluations of Micromedex Solutions at Truven Health, or The American Hospital Formulary Service Drug Information).

### REFERENCES

1. Neulasta [package insert]. Thousand Oaks, CA: Amgen Inc.; January 2025.
2. Armlupeg [package insert]. Naples, FL: Lupin Pharmaceuticals, Inc.; December 2025.
3. Ennumo [package insert]. Raleigh, NC: Accord BioPharma Inc.; May 2026.
4. Fulphila [package insert]. Cambridge, MA: Biocon Biologics, Inc.; June 2023.
5. Udenyca [package insert]. Redwood City, CA: Coherus BioSciences, Inc.; December 2023.
6. Ziextenzo [package insert]. Princeton, NJ: Sandoz Inc.; February 2024.
7. Nyvepria [package insert]. Lake Forest, IL: Hospira, Inc.; March 2023.
8. Fylnetra [package insert]. Piscataway, NJ: Kashiv BioSciences, LLC; April 2025.
9. Stimufend [package insert]. Lake Zurich, IL: Fresenius Kabi USA, LLC; September 2023.
10. The NCCN Drugs & Biologics Compendium © 2025 National Comprehensive Cancer Network, Inc. Available at: <https://www.nccn.org> Accessed June 12, 2025.
11. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Hematopoietic Growth Factors. Version 1.2025. [https://www.nccn.org/professionals/physician\\_gls/pdf/growthfactors.pdf](https://www.nccn.org/professionals/physician_gls/pdf/growthfactors.pdf) Accessed June 19, 2025.
12. IBM Micromedex® DRUGDEX ® (electronic version). IBM Watson Health, Greenwood Village, Colorado, USA. Available at <https://www.micromedexsolutions.com> (Accessed: June 19, 2025).
13. Smith TJ, Bohlke K, Lyman GH, et al. Recommendations for the use of white blood cell growth factors: American Society of Clinical Oncology Clinical Practice Guideline Update. *J Clin Oncol*. 2015;33(28):3199-3212.
14. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Hairy Cell Leukemia. Version 1.2025. [https://www.nccn.org/professionals/physician\\_gls/pdf/hairy\\_cell.pdf](https://www.nccn.org/professionals/physician_gls/pdf/hairy_cell.pdf) Accessed June 19, 2025.
15. Smith TJ, Khatcheressian J, Lyman GH, et al. 2006 update of recommendations for the use of white blood cell growth factors: an evidence-based clinical practice guideline. *J Clin Oncol*. 2006;24(19):3187-3205.
16. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Gestational Trophoblastic Neoplasia. Version 3.2025. [https://www.nccn.org/professionals/physician\\_gls/pdf/gtn.pdf](https://www.nccn.org/professionals/physician_gls/pdf/gtn.pdf) Accessed June 19, 2025.
17. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology: Wilms Tumor (Nephroblastoma). Version 2.2025. [https://www.nccn.org/professionals/physician\\_gls/pdf/wilms\\_tumor.pdf](https://www.nccn.org/professionals/physician_gls/pdf/wilms_tumor.pdf) Accessed June 19, 2025.

### EFFECTIVE DATE

ID\_CHS\_19312025b\_1026