



Medical Policy Manual **Approved Rev: Do Not Implement until 6/30/26**

Rituximab Products (Rituxan®, Rituximab-abbs [Truxima®], Rituximab-arrx [Riabni™] and Rituximab-pvvr [Ruxience®])

Treatment of Rheumatoid Arthritis and Other Conditions (Non-Oncology Indications)

Some agents on this policy may require step therapy See “Step Therapy Requirements for Provider Administered Specialty Medications” Document at:

https://www.bcbst.com/docs/providers/Comm_BC_PAD_Step_Therapy_Guide.pdf

IMPORTANT REMINDER

We develop Medical Policies to provide guidance to Members and Providers. This Medical Policy relates only to the services or supplies described in it. The existence of a Medical Policy is not an authorization, certification, explanation of benefits or a contract for the service (or supply) that is referenced in the Medical Policy. For a determination of the benefits that a Member is entitled to receive under his or her health plan, the Member's health plan must be reviewed. If there is a conflict between the medical policy and a health plan or government program (e.g., TennCare), the express terms of the health plan or government program will govern.

POLICY

INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

- Granulomatosis with Polyangiitis (GPA) (Wegener's Granulomatosis) and Microscopic Polyangiitis (MPA) in adult and pediatric patients 2 years of age and older* in combination with glucocorticoids (*pediatric indication applies to Rituxan only).
- Rheumatoid Arthritis (RA) in combination with methotrexate in adult patients with moderately-to severely-active RA who have inadequate response to one or more TNF antagonist therapies.
- Non-Hodgkin's lymphoma (NHL):
(Not addressed in this policy – Refer to Rituxan-Ruxience-Truxima-Riabni-Oncology Medical Policy)
- Chronic lymphocytic leukemia (CLL):
(Not addressed in this policy – Refer to Rituxan-Ruxience-Truxima-Riabni-Oncology Medical Policy)
- Moderate to severe pemphigus vulgaris in adult patients.
- Mature B-cell acute leukemia (B-AL):
(Not addressed in this policy - Refer to Rituxan-Ruxience-Truxima-Riabni Oncology Medical Policy)

Compendial Uses

- Sjögren's syndrome
- Multiple sclerosis, relapsing-remitting
- Neuromyelitis optica (i.e., neuromyelitis optica spectrum disorder, NMOSD, Devic disease)
- Autoimmune blistering disease
- Cryoglobulinemia
- Solid organ transplant
- Opsoclonus-myoclonus ataxia



Medical Policy Manual **Approved Rev: Do Not Implement until 6/30/26**

- Systemic lupus erythematosus (SLE)
- Myasthenia gravis, refractory
- Membranous nephropathy
- Eosinophilic granulomatosis with polyangiitis (EGPA)
- Susac syndrome
- For other compendial uses, refer to Rituxan-Ruxience-Truxima-Riabni-Oncology Medical Policy

All other indications are considered experimental/investigational and not medically necessary.

DOCUMENTATION

Submission of the following information is necessary to initiate the prior authorization review:

Rheumatoid arthritis (RA)

Initial requests

- Chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.
- Laboratory results, chart notes, or medical record documentation of biomarker testing (i.e., rheumatoid factor [RF], anti-cyclic citrullinated peptide [anti-CCP], and C-reactive protein [CRP] and/or erythrocyte sedimentation rate [ESR]) (if applicable).

Continuation requests

Chart notes or medical record documentation supporting positive clinical response.

Sjögren's syndrome, cryoglobulinemia, opsoclonus-myoclonus-ataxia, and systemic lupus erythematosus (initial requests only)

Chart notes, medical record documentation, or claims history supporting previous medications tried, including response to therapy.

PRESCRIBER SPECIALITIES

This medication must be prescribed by or in consultation with one of the following:

- RA, GPA (Wegener's granulomatosis), MPA, EGPA, pauci-immune glomerulonephritis, SLE: rheumatologist, immunologist, nephrologist
- Sjogren's syndrome: rheumatologist, ophthalmologist, immunologist
- Multiple sclerosis, NMOSD, myasthenia gravis, opsoclonus-myoclonus-ataxia, Susac syndrome: neurologist, immunologist, rheumatologist
- Autoimmune blistering disease: dermatologist, immunologist
- Cryoglobulinemia: hematologist, rheumatologist, neurologist, nephrologist
- Solid organ transplant: immunologist, transplant specialist
- Membranous nephropathy: nephrologist, rheumatologist

EXCLUSIONS

- Member will not receive Rituxan, Ruxience, Truxima, or Riabni with any other biologic drug or targeted synthetic drug for RA.
- Member will not receive Rituxan, Ruxience, Truxima, or Riabni with other disease modifying multiple sclerosis agents (Note: Ampyra and Nuedexta are not disease modifying).



Medical Policy Manual **Approved Rev: Do Not Implement until 6/30/26**

- Member will not use Rituxan, Ruxience, Truxima, or Riabni concomitantly with other biologics for the treatment of neuromyelitis optica.

COVERAGE CRITERIA

Rheumatoid arthritis (RA)

Authorization of 12 months may be granted for adults who have previously received a biologic or targeted synthetic drug (e.g., Rinvoq, Xeljanz) indicated for the treatment of moderately to severely active rheumatoid arthritis. The requested medication must be prescribed in combination with methotrexate (MTX) or leflunomide unless the member has a clinical reason not to use methotrexate or leflunomide (see Appendix).

Authorization of 12 months may be granted for treatment of adults with moderately to severely active RA when all of the following criteria are met:

- The member meets either of the following criteria:
 - The member has been tested for either of the following biomarkers and the test was positive:
 - Rheumatoid factor (RF)
 - Anti-cyclic citrullinated peptide (anti-CCP)
 - The member has been tested for ALL of the following biomarkers:
 - RF
 - Anti-CCP
 - C-reactive protein (CRP) and/or erythrocyte sedimentation rate (ESR)
- The member is prescribed the requested medication in combination with methotrexate or leflunomide or has a clinical reason not to use methotrexate or leflunomide (see Appendix).
- The member meets either of the following criteria:
 - The member has experienced an inadequate response to at least a 3-month trial of MTX despite adequate dosing (i.e., titrated to at least 15 mg/week)
 - The member had an intolerable adverse effect or contraindication to MTX (see Appendix), and an inadequate response to another conventional drug (e.g., hydroxychloroquine, leflunomide, sulfasalazine).

Granulomatosis with polyangiitis (GPA) (Wegener's granulomatosis), microscopic polyangiitis (MPA), eosinophilic granulomatosis with polyangiitis (EGPA), and pauci-immune glomerulonephritis

Authorization of 12 months may be granted for treatment of GPA, MPA, EGPA, or pauci-immune glomerulonephritis.

Sjögren's syndrome

Authorization of 12 months may be granted for treatment of Sjögren's syndrome when corticosteroids and other immunosuppressive agents were ineffective.

Multiple sclerosis

Authorization of 12 months may be granted for treatment of relapsing-remitting multiple sclerosis (RRMS).

Neuromyelitis optica (i.e., neuromyelitis optica spectrum disorder [NMOSD], Devic disease)

Authorization of 12 months may be granted for treatment of neuromyelitis optica (i.e., neuromyelitis optica spectrum disorder [NMOSD], Devic disease).

Medical Policy Manual **Approved Rev: Do Not Implement until 6/30/26**

Autoimmune blistering disease

Authorization of 12 months may be granted for treatment of autoimmune blistering disease (e.g., pemphigus vulgaris, pemphigus foliaceus, bullous pemphigoid, cicatricial pemphigoid, epidermolysis bullosa acquisita and paraneoplastic pemphigus).

Cryoglobulinemia

Authorization of 12 months may be granted for treatment of cryoglobulinemia when corticosteroids and other immunosuppressive agents were ineffective.

Solid organ transplant

Authorization of 3 months may be granted for treatment of solid organ transplant and prevention of antibody-mediated rejection in solid organ transplant.

Opsoclonus-myoclonus-ataxia

Authorization of 12 months may be granted for treatment of opsoclonus-myoclonus-ataxia associated with neuroblastoma when the member is refractory to steroids and chemotherapy.

Systemic lupus erythematosus

Authorization of 12 months may be granted for the treatment of systemic lupus erythematosus that is refractory to immunosuppressive therapy.

Myasthenia gravis

Authorization of 12 months may be granted for treatment of refractory myasthenia gravis.

Membranous nephropathy

Authorization of 12 months may be granted for treatment of membranous nephropathy when the member is at moderate to high risk for disease progression.

Susac syndrome

Authorization of 12 months may be granted for treatment of Susac Syndrome.

CONTINUATION OF THERAPY

Rheumatoid arthritis

Authorization of 12 months may be granted for continued treatment in all adult members (including new members) who are using the requested medication for moderately to severely active rheumatoid arthritis and who achieve or maintain a positive clinical response after at least two doses of therapy with Rituxan, Ruxience, Truxima, or Riabni as evidenced by disease activity improvement of at least 20% from baseline in tender joint count, swollen joint count, pain, or disability.

Multiple sclerosis



Medical Policy Manual **Approved Rev: Do Not Implement until 6/30/26**

Authorization of 12 months may be granted for continued treatment in members requesting reauthorization for relapsing-remitting multiple sclerosis who are experiencing disease stability or improvement while receiving Rituxan, Ruxience, Truxima, or Riabni.

Other indications

Authorization of 12 months may be granted for continued treatment in all members (including new members) requesting reauthorization who are receiving benefit from therapy.

APPENDIX

Examples of clinical reasons to avoid pharmacologic treatment with methotrexate or leflunomide

- Clinical diagnosis of alcohol use disorder, alcoholic liver disease or other chronic liver disease
- Drug interaction
- Risk of treatment-related toxicity
- Pregnancy or currently planning pregnancy
- Breastfeeding
- Significant comorbidity prohibits use of systemic agents (e.g., liver or kidney disease, blood dyscrasias, uncontrolled hypertension)
- Hypersensitivity
- History of intolerance or adverse event

MEDICATION QUANTITY LIMITS

Drug Name	Diagnosis	Maximum Dosing Regimen
Rituxan (Rituximab)	Cryoglobulinemia	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every weeks
Rituxan (Rituximab)	Granulomatosis with Polyangiitis (GPA) (Wegener's Granulomatosis) and Microscopic Polyangiitis (MPA)	Route of Administration: Intravenous <u>≥18 year(s)</u> 375mg/m ² every week for 4 weeks, followed by 500 mg every 2 weeks for 2 doses (start no sooner than 16 to 24 weeks after last induction dose), followed by 500 mg every 6 months based on clinical evaluation 1000mg every 2 weeks for 2 doses, followed by 1000 mg every 4 to 6 months <u>≥2 to <18 year(s)</u> 375mg/m ² every week for 4 weeks, followed by 250 mg/m ² every 2 weeks for 2 doses (start no sooner than 16 to 24 weeks after last induction dose), followed by 250 mg/m ² every 6 months based on clinical evaluation



Medical Policy Manual **Approved Rev: Do Not Implement until 6/30/26**

		750mg/m ² (max 1000 mg/dose) every 2 weeks for 2 doses, followed by 750 mg/m ² (max 1000 mg/dose) every 4 to 6 months
Rituxan (Rituximab)	Membranous Nephropathy	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Rituxan (Rituximab)	Multiple Sclerosis	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Rituxan (Rituximab)	Myasthenia Gravis	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Rituxan (Rituximab)	Neuromyelitis Optica (i.e. Neuromyelitis Optica Spectrum Disorder, NMOSD, or Devic Disease)	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Rituxan (Rituximab)	Opsoclonus-Myoclonus Ataxia	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Rituxan (Rituximab)	Pemphigus Vulgaris (PV) and other Autoimmune Blistering Disease (e.g., Pemphigus Foliaceus, Bullous Pemphigoid, Cicatricial Pemphigoid, Epidermolysis Bullosa Acquisita and Paraneoplastic Pemphigus)	Route of Administration: Intravenous ≥ 18 year(s) 375mg/m ² every week for 4 doses per treatment course; may repeat treatment after at least 6 months if necessary Initial: 1000mg every 2 weeks for 2 doses Maintenance: 500mg at month 12 and every 6 months thereafter 1000mg on relapse; subsequent infusions no sooner than 16 weeks following previous infusion
Rituxan (Rituximab)	Rheumatoid Arthritis	Route of Administration: Intravenous ≥ 18 year(s) 1000mg x 2 doses given 2 weeks apart (one course); Subsequent courses may be given every 24 weeks or based on clinical evaluation, but not sooner than every 16 weeks
Rituxan (Rituximab)	Sjögren's Syndrome	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Rituxan (Rituximab)	Solid Organ Transplant	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Rituxan (Rituximab)	Systemic Lupus Erythematosus	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every weeks
Truxima (Rituximab-abbs)	Cryoglobulinemia	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week



Medical Policy Manual **Approved Rev: Do Not Implement until 6/30/26**

Truxima (Rituximab-abbs)	Granulomatosis with Polyangiitis (GPA) (Wegener's Granulomatosis) and Microscopic Polyangiitis (MPA)	Route of Administration: Intravenous ≥18 year(s) 375mg/m ² every week for 4 weeks, followed by 500 mg every 2 weeks for 2 doses (start no sooner than 16 to 24 weeks after last induction dose), followed by 500 mg every 6 months based on clinical evaluation 1000mg every 2 weeks for 2 doses, followed by 1000 mg every 4 to 6 months
Truxima (Rituximab-abbs)	Membranous Nephropathy	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Truxima (Rituximab-abbs)	Multiple Sclerosis	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Truxima (Rituximab-abbs)	Myasthenia Gravis	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Truxima (Rituximab-abbs)	Neuromyelitis Optica (i.e. Neuromyelitis Optica Spectrum Disorder, NMOSD, or Devic Disease)	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Truxima (Rituximab-abbs)	Opsoclonus-Myoclonus Ataxia	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Truxima (Rituximab-abbs)	Pemphigus Vulgaris (PV) and other Autoimmune Blistering Disease (e.g., Pemphigus Foliaceus, Bullous Pemphigoid, Cicatricial Pemphigoid, Epidermolysis Bullosa Acquisita and Paraneoplastic Pemphigus)	Route of Administration: Intravenous ≥18 year(s) 375mg/m ² every week for 4 doses per treatment course; may repeat treatment after at least 6 months if necessary Initial: 1000mg every 2 weeks for 2 doses Maintenance: 500mg at month 12 and every 6 months thereafter 1000mg on relapse; subsequent infusions no sooner than 16 weeks following previous infusion
Truxima (Rituximab-abbs)	Rheumatoid Arthritis	Route of Administration: Intravenous ≥18 year(s) 1000mg x 2 doses given 2 weeks apart (one course); Subsequent courses may be given every 24 weeks or based on clinical evaluation, but not sooner than every 16 weeks
Truxima (Rituximab-abbs)	Sjögren's Syndrome	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)



Medical Policy Manual **Approved Rev: Do Not Implement until 6/30/26**

Truxima (Rituximab-abbs)	Solid Organ Transplant	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Truxima (Rituximab-abbs)	Systemic Lupus Erythematosus	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Riabni (Rituximab-arxx)	Cryoglobulinemia	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Riabni (Rituximab-arxx)	Granulomatosis with Polyangiitis (GPA) (Wegener's Granulomatosis) and Microscopic Polyangiitis (MPA)	Route of Administration: Intravenous ≥18 year(s) 375mg/m ² every week for 4 weeks, followed by 500 mg every 2 weeks for 2 doses (start no sooner than 16 to 24 weeks after last induction dose), followed by 500 mg every 6 months based on clinical evaluation 1000mg every 2 weeks for 2 doses, followed by 1000 mg every 4 to 6 months
Riabni (Rituximab-arxx)	Membranous Nephropathy	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Riabni (Rituximab-arxx)	Multiple Sclerosis	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Riabni (Rituximab-arxx)	Myasthenia Gravis	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Riabni (Rituximab-arxx)	Neuromyelitis Optica (i.e. Neuromyelitis Optica Spectrum Disorder, NMOSD, or Devic Disease)	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Riabni (Rituximab-arxx)	Opsoclonus-Myoclonus Ataxia	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Riabni (Rituximab-arxx)	Pemphigus Vulgaris (PV) and other Autoimmune Blistering Disease (e.g., Pemphigus Foliaceus, Bullous Pemphigoid, Cicatricial Pemphigoid, Epidermolysis Bullosa Acquisita and Paraneoplastic Pemphigus)	Route of Administration: Intravenous ≥18 year(s) 375mg/m ² every week for 4 doses per treatment course; may repeat treatment after at least 6 months if necessary Initial: 1000mg every 2 weeks for 2 doses Maintenance: 500mg at month 12 and every 6 months thereafter 1000mg on relapse; subsequent infusions no sooner than 16 weeks following previous infusion
Riabni (Rituximab-arxx)	Rheumatoid Arthritis	Route of Administration: Intravenous ≥18 year(s) 1000mg x 2 doses given 2 weeks apart



Medical Policy Manual **Approved Rev: Do Not Implement until 6/30/26**

		(one course); Subsequent courses may be given every 24 weeks or based on clinical evaluation, but not sooner than every 16 weeks
Riabni (Rituximab-arrx)	Sjögren's Syndrome	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Riabni (Rituximab-arrx)	Solid Organ Transplant	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Riabni (Rituximab-arrx)	Systemic Lupus Erythematosus	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Ruxience (Rituximab-pvvr)	Cryoglobulinemia	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Ruxience (Rituximab-pvvr)	Granulomatosis with Polyangiitis (GPA) (Wegener's Granulomatosis) and Microscopic Polyangiitis (MPA)	Route of Administration: Intravenous ≥18 year(s) 375mg/m ² every week for 4 weeks, followed by 500 mg every 2 weeks for 2 doses (start no sooner than 16 to 24 weeks after last induction dose), followed by 500 mg every 6 months based on clinical evaluation 1000mg every 2 weeks for 2 doses, followed by 1000 mg every 4 to 6 months
Ruxience (Rituximab-pvvr)	Membranous Nephropathy	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Ruxience (Rituximab-pvvr)	Multiple Sclerosis	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Ruxience (Rituximab-pvvr)	Myasthenia Gravis	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Ruxience (Rituximab-pvvr)	Neuromyelitis Optica (i.e. Neuromyelitis Optica Spectrum Disorder, NMOSD, or Devic Disease)	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week
Ruxience (Rituximab-pvvr)	Opsoclonus-Myoclonus Ataxia	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Ruxience (Rituximab-pvvr)	Pemphigus Vulgaris (PV) and other Autoimmune Blistering Disease (e.g., Pemphigus Foliaceus, Bullous Pemphigoid, Cicatricial Pemphigoid, Epidermolysis Bullosa Acquisita and Paraneoplastic Pemphigus)	Route of Administration: Intravenous ≥18 year(s) 375mg/m ² every week for 4 doses per treatment course; may repeat treatment after at least 6 months if necessary Initial: 1000mg every 2 weeks for 2 doses Maintenance: 500mg at month 12 and every 6 months thereafter

Medical Policy Manual **Approved Rev: Do Not Implement until 6/30/26**

		1000mg on relapse; subsequent infusions no sooner than 16 weeks following previous infusion
Ruxience (Rituximab-pvvr)	Rheumatoid Arthritis	Route of Administration: Intravenous ≥18 year(s) 1000mg x 2 doses given 2 weeks apart (one course); Subsequent courses may be given every 24 weeks or based on clinical evaluation, but not sooner than every 16 weeks
Ruxience (Rituximab-pvvr)	Sjögren's Syndrome	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Ruxience (Rituximab-pvvr)	Solid Organ Transplant	Route of Administration: Intravenous 1000mg (frequency should not be more frequent than every 14 days)
Ruxience (Rituximab-pvvr)	Systemic Lupus Erythematosus	Route of Administration: Intravenous 1000mg every 2 weeks or 375 mg/m ² every week

APPLICABLE TENNESSEE STATE MANDATE REQUIREMENTS

BlueCross BlueShield of Tennessee's Medical Policy complies with Tennessee Code Annotated Section 56-7-2352 regarding coverage of off-label indications of Food and Drug Administration (FDA) approved drugs when the off-label use is recognized in one of the statutorily recognized standard reference compendia or in the published peer-reviewed medical literature.

ADDITIONAL INFORMATION

For appropriate chemotherapy regimens, dosage information, contraindications, precautions, warnings, and monitoring information, please refer to one of the standard reference compendia (e.g., the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) published by the National Comprehensive Cancer Network®, Drugdex Evaluations of Micromedex Solutions at Truven Health, or The American Hospital Formulary Service Drug Information).

REFERENCES

1. Rituxan [package insert]. South San Francisco, CA: Genentech, Inc.; December 2021.
2. Truxima [package insert]. North Wales, PA: Teva Pharmaceuticals USA, Inc.; June 2025.
3. Methotrexate [package insert]. Cranbury, NJ: Sun Pharmaceutical Industries, Inc.; August 2021.
4. IBM Micromedex [Internet database]. Ann Arbor, MI: Truven Health Analytics. Updated periodically. Accessed April 8, 2025.
5. Dass S, Bowman SJ, Vital EM, et al. Reduction of fatigue in Sjögren syndrome with rituximab: results of a double blind, placebo-controlled study. *Ann Rheum Dis.* 2008;67:1541-1544.
6. Meijer JM, Meiners PM, Vissink A, et al. Effectiveness of rituximab treatment in primary Sjögren's syndrome: a randomized, double-blind, placebo-controlled trial. *Arthritis Rheum.* 2010;62(4):960-8.
7. Smolen JS, Landewé R, Billsma J, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Ann Rheum Dis.* 2020;79:685-699.

Medical Policy Manual **Approved Rev: Do Not Implement until 6/30/26**

8. Singh JA, Saag KG, Bridges SL Jr, et al. 2015 American -College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Rheumatol*. 2016;68(1):1-26.
9. Saag KG, Teng GG, Patkar NM, et al. American College of Rheumatology 2008 recommendations for the use of nonbiologic and biologic disease-modifying antirheumatic drugs in rheumatoid arthritis. *Arthritis Rheum*. 2008;59(6):762-784.
10. Hauser SL, Waubant E, Arnold DL, et al. B-cell depletion with rituximab in relapsing-remitting multiple sclerosis. *N Engl J Med*. 2008;358:676-688.
11. Scott, T.F., Frohman, E.M., DeSeze, J., (2011). Evidence-based guideline: Clinical evaluation and treatment of transverse myelitis: Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *American Academy of Neurology*. 77: 2128-2134.
12. Kumpfel, T., Gighuber, K., et al. (2023). Update on the diagnosis and treatment of neuromyelitis optica spectrum disorders (NMOSD)-revised recommendations of the Neuromyelitis Optica Study Group (NEMOS). Part II: Attack therapy and long-term management. *J Neurol* 271: 141-176.
13. De Vita S, Quartuccio L, Isola M, et al. A randomized controlled trial of rituximab for the treatment of severe cryoglobulinemic vasculitis. *Arthritis Rheum*. 2012;64(3):843-53.
14. Sneller MC, Hu Z, Langford CA. A randomized controlled trial of rituximab following failure of antiviral therapy for hepatitis C virus-associated cryoglobulinemic vasculitis. *Arthritis Rheum*. 2012 Mar; 64(3):835-42.
15. Terrier B, Krastinova E, Marie I, et al. Management of noninfectious mixed cryoglobulinemia vasculitis: data from 242 cases included in the CryoVas survey. *Blood*. 2012 Jun 21; 119(25):5996-6004.
16. Trappe R, Oertel S, Leblond V, et al. Sequential treatment with rituximab followed by CHOP chemotherapy in adult B-cell post-transplant lymphoproliferative disorder (PTLD): the prospective international multicentre phase 2 PTL-1 trial. *Lancet Oncol* 2012 Feb; 13 (2):196-206.
17. The American Society of Transplantation Infectious Diseases Guidelines. *Am J Transplant* 2009; 9 (Suppl 4):S92.
18. Bell J, Moran C, Blatt J. Response to rituximab in a child with neuroblastoma and opsoclonus-myoclonus. *Pediatr Blood Cancer* 2008; 50:370.
19. Hertl M, Geller S. Initial management of pemphigus vulgaris and pemphigus foliaceus. UpToDate [online serial]. Waltham, MA: UpToDate; reviewed August 2021.
20. American Society of Health System Pharmacists. AHFS DI. Bethesda, MD. Electronic version, 2019. Available with subscription. URL: <http://online.lexi.com/lco/action/home>. Accessed April 8, 2025.
21. DRUGDEX System (electronic version). Micromedex Truven Health Analytics. Available with subscription. URL: www.micromedexsolutions.com. Accessed April 8, 2025.
22. Ruxience [package insert]. NY, NY: Pfizer Biosimilars; June 2025.
23. Riabni [package insert]. Thousand Oaks, CA: Amgen Inc.; June 2025.
24. Aletaha D, Neogi T, Silman, et al. 2010 Rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Arthritis Rheum*. 2010;62(9):2569-81.
25. Smolen JS, Aletaha D. Assessment of rheumatoid arthritis activity in clinical trials and clinical practice. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. Available with subscription. URL: www.uptodate.com. Accessed November 7, 2022.
26. Murrell DF, Peña S, Joly P, et al. Diagnosis and management of pemphigus: Recommendations of an international panel of experts. *J Am Acad Dermatol*. 2020;82(3):575-585.e1.
27. Joly P, Horvath B, Patsatsi A, et al. Updated S2K guidelines on the management of pemphigus vulgaris and foliaceus initiated by the European Academy of Dermatology and Venereology (EADV). *J Eur Acad Dermatol Venereol*. 2020;34(9):1900-1913.
28. Micromedex Solutions [database online]. Truven Health Analytics, Greenwood Village, Colorado, USA. Available at: <http://www.micromedexsolutions.com/>. Accessed March 27, 2025.
29. Fraenkel L, Bathon JM, England BR, et al. 2021 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. *Arthritis Care Res*. 2021;0:1-16.
30. Menditto VG, Rossetti G, Olivari D, et al. Rituximab for eosinophilic granulomatosis with polyangiitis: a systematic review of observational studies. *Rheumatology (Oxford)*. 2021;60(4):1640-1650.



Medical Policy Manual **Approved Rev: Do Not Implement until 6/30/26**

31. Chung SA, Langford CA, Maz M, et al. 2021 American College of Rheumatology/Vasculitis Foundation Guideline for the Management of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. *Arthritis Rheumatol.* 2021;73(8):1366-1383.
32. Poupart J, Giovannelli J, Deschamps R, Audoin B, Ciron J, Maillart E, Papeix C, Collongues N, Bourre B, Cohen M, Wiertelowski S, Outteryck O, Laplaud D, Vukusic S, Marignier R, Zephir H; NOMADMUS study group. Evaluation of efficacy and tolerability of first-line therapies in NMOSD. *Neurology.* 2020 Apr 14;94(15):e1645-e1656.
33. Kidney Disease: Improving Global Outcomes (KDIGO) Glomerular Diseases Work Group. KDIGO 2021 Clinical Practice Guideline for the Management of Glomerular Diseases. *Kidney Int.* 2021 Oct;100(4S):S1-S276.
34. Kidney Disease: Improving Global Outcomes (KDIGO) Glomerular Diseases Work Group, KDIGO 2024 Clinical Practice Guideline for the Management of Antineutrophil Cytoplasmic Antibody (ANCA)-Associated Vasculitis. *Kidney Int.* 2024;105(Suppl 3S):S71–S116.
35. Rennebohm RM, Asdaghi N, Srivastava S, et al. Guidelines for treatment of Susac Syndrome – An update. *International Journal of Stroke.* 2018;0(0):1-11.
36. Rae-Grant A, Day GS, Marrier RA, et al., Practice guideline recommendations summary: Disease-modifying therapies for adults with multiple sclerosis. *Neurology.* 2018;90(17):777-788.

EFFECTIVE DATE 6/30/2026

ID_CHS_2025a