



## Medical Policy Manual **Approved Rev: Do Not Implement until 7/31/26**

### Evinacumab-dgnb (Evkeeza™)

#### IMPORTANT REMINDER

We develop Medical Policies to provide guidance to Members and Providers. This Medical Policy relates only to the services or supplies described in it. The existence of a Medical Policy is not an authorization, certification, explanation of benefits or a contract for the service (or supply) that is referenced in the Medical Policy. For a determination of the benefits that a Member is entitled to receive under his or her health plan, the Member's health plan must be reviewed. If there is a conflict between the medical policy and a health plan or government program (e.g., TennCare), the express terms of the health plan or government program will govern.

#### POLICY

#### INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

##### FDA-Approved Indications

Evkeeza is indicated as an adjunct to diet and exercise and other low-density lipoprotein-cholesterol (LDL-C) lowering therapies to reduce LDL-C in adult and pediatric patients, aged 1 year and older, with homozygous familial hypercholesterolemia (HoFH).

All other indications are considered experimental/investigational and not medically necessary.

#### DOCUMENTATION

Submission of the following information is necessary to initiate the prior authorization review:

##### Initial Requests:

- Genetic testing, **chart notes**, or medical record **documentation** confirming the diagnosis of HoFH.
- With clinical atherosclerotic cardiovascular disease (ASCVD): Chart notes confirming clinical ASCVD (if applicable) (see Appendix).
- **Current** LDL-C level dated within the six months preceding the authorization request.
- For members 10 years of age and older: chart notes, medical record documentation, or claims history confirming the member is currently on maximally tolerated lipid-lowering therapy.
- For members 7 years of age to less than 10 years of age: chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.

##### Continuation Requests:

- **Current** LDL-C level dated within the six months preceding the authorization request.
- For members 10 years of age and older: **chart notes, medical record documentation, or claims history confirming the member is currently on maximally tolerated lipid-lowering therapy.**
- For members 7 years of age to less than 10 years of age: **chart notes, medical record documentation, or claims history supporting previous medications tried (if applicable), including response to therapy. If therapy is not advisable, documentation of clinical reason to avoid therapy.**

#### PRESCRIBER SPECIALTIES

## Medical Policy Manual **Approved Rev: Do Not Implement until 7/31/26**

This medication must be prescribed by or in consultation with a cardiologist, endocrinologist, lipid specialist, geneticist, or a prescriber specialized in the treatment of HoFH.

### COVERAGE CRITERIA

#### Homozygous Familial Hypercholesterolemia (HoFH)

Authorization of 6 months may be granted for members 1 year of age or older for treatment of homozygous familial hypercholesterolemia (HoFH) when all of the following criteria are met:

- Member has a documented diagnosis of HoFH confirmed by any of the following criteria:
  - Variant in two low-density lipoprotein receptor (LDLR) alleles.
  - Presence of homozygous or compound heterozygous variants in apolipoprotein B (APOB) or proprotein convertase subtilisin-kexin type 9 (PCSK9) gene.
  - Member has compound heterozygosity or homozygosity for variants in the gene encoding low-density lipoprotein receptor adaptor protein 1 (LDLRAP1).
  - Member has an untreated LDL-C of > 400 mg/dL and has either of the following:
    - Presence of cutaneous or tendinous xanthomas before the age of 10 years.
    - An untreated LDL-C level of  $\geq 190$  mg/dL in both parents.
- Prior to initiation of treatment with the requested medication, member meets/ has met either of the following criteria:
  - Member has a treated LDL-C level  $\geq 70$  mg/dL
  - Member has a treated LDL-C level  $\geq 55$  mg/dL and meets either of the following criteria:
    - Member has a history of a clinical ASCVD event (see Appendix).
    - Member has major ASCVD risk factors (e.g., 65 years of age or older, familial hypercholesterolemia, diabetes, chronic kidney disease, history of congestive heart failure).
- Prior to initiation of treatment with the requested medication, member meets/has met one of the following criteria:
  - Member is 10 years of age or older and meets both of the following criteria:
    - Member is receiving stable treatment with at least 3 lipid-lowering therapies (e.g., statins, ezetimibe, PCSK9 directed therapy) at the maximally tolerated dose.
    - Member will continue to receive concomitant lipid-lowering therapy at the maximally tolerated dose.
  - Member is 7 years of age to less than 10 years of age and meets either of the following criteria:
    - Member is receiving stable treatment with at least one lipid-lowering therapy (e.g., statins, LDL apheresis) at the maximally tolerated dose and will continue to receive concomitant lipid-lowering therapy at the maximally tolerated dose.
    - Member has an intolerance or contraindication to other lipid-lowering therapies.
  - Member is 1 year of age to less than 7 years of age.

### CONTINUATION OF THERAPY

Authorization of 12 months may be granted for continued treatment in members (including new members) who meet all of the following criteria:

- Member meets one of the following criteria:
  - Member is 10 years of age or older and is currently receiving concomitant lipid-lowering therapy at the maximally tolerated dose.
  - Member is 7 years of age to less than 10 years of age and meets either of the following criteria:
    - Member is currently receiving concomitant lipid-lowering therapy at the maximally tolerated dose.
    - Member has an intolerance or contraindication to other lipid-lowering therapies.
  - Member is 1 year of age to less than 7 years of age.

## Medical Policy Manual **Approved Rev: Do Not Implement until 7/31/26**

- The member is receiving benefit from therapy. Benefit is defined as either of the following:
  - LDL-C is now at goal.
  - Member has had at least 30% reduction of LDL-C from baseline.

### APPENDIX

#### Clinical ASCVD

- Acute coronary syndromes
- Myocardial infarction
- Stable or unstable angina
- Coronary or other arterial revascularization procedure (e.g., percutaneous coronary intervention [PCI], coronary artery bypass graft [CABG] surgery)
- Stroke of presumed atherosclerotic origin
- Transient ischemic attack (TIA)
- Non-cardiac peripheral arterial disease (PAD) of presumed atherosclerotic origin (e.g., carotid artery stenosis, lower extremity PAD)
- Obstructive coronary artery disease (defined as  $\geq 50\%$  stenosis on cardiac computed tomography angiogram or catheterization)
- Coronary artery calcium (CAC) Score  $\geq 300$

### APPLICABLE TENNESSEE STATE MANDATE REQUIREMENTS

BlueCross BlueShield of Tennessee's Medical Policy complies with Tennessee Code Annotated Section 56-7-2352 regarding coverage of off-label indications of Food and Drug Administration (FDA) approved drugs when the off-label use is recognized in one of the statutorily recognized standard reference compendia or in the published peer-reviewed medical literature.

### ADDITIONAL INFORMATION

For appropriate chemotherapy regimens, dosage information, contraindications, precautions, warnings, and monitoring information, please refer to one of the standard reference compendia (e.g., the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) published by the National Comprehensive Cancer Network®, Drugdex Evaluations of Micromedex Solutions at Truven Health, or The American Hospital Formulary Service Drug Information).

### REFERENCES

1. Evkeeza [package insert]. Tarrytown, NY: Regeneron Pharmaceuticals Inc.; September 2025.
2. Raal FJ, Rosenson RS, Reeskamp LF, et al. Evinacumab for homozygous familial hypercholesterolemia. *N Engl J Med.* 2020;383(8):711-720.
3. Mach F, Koskinas KC, Roeters van Lennep JE, et al. 2025 Focused update of the 2019 ECS/EAS guidelines for the management of dyslipidaemias. *Eur Heart J.* 2025; 46: 4359-4378.
4. Cuchel M, Raal FJ, Hegele RA, et al. 2023 Update on European Atherosclerosis Society consensus statement on homozygous familial hypercholesterolaemia: new treatments and clinical guidance. *Eur Heart J.* 2023;44(25):2277-2291.
5. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: a report of the American College of



## Medical Policy Manual **Approved Rev: Do Not Implement until 7/31/26**

Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. Circulation. 2019;139(25):e1082– e1143.

6. Wiegman A, Greber-Platzer S, Ali S, et al. Evinacumab in pediatric patients with homozygous familial hypercholesterolemia. Circulation. 2024;149(5):343-353.
7. Lloyd-Jones DM, Morris PB, Ballantyne CM, et al. 2022 ACC expert consensus decision pathway on the role of nonstatin therapies for LDL-cholesterol lowering in the management of atherosclerotic cardiovascular disease risk: A report of the American College of Cardiology Solution Set Oversight Committee. J Am Coll Cardiol. 2022;80(14):1366–1418.
8. Jacobson TA, Ito MK, Maki KC, et al. National Lipid Association recommendations for patient-centered management of dyslipidemia: part 1 — full report. J Clin Lipidol. 2015;9:129–169.
9. Min JK, Labounty TM, Gomez MJ, et al. Incremental prognostic value of coronary computed tomographic angiography over coronary artery calcium score for risk prediction of major adverse cardiac events in asymptomatic diabetic individuals. Atherosclerosis. 2014;232(2):298-304.
10. Budoff MJ, Kinnering A, Gransar H, et al. When does a calcium score equate to secondary prevention?: Insights from the multinational CONFIRM registry. JACC Cardiovasc Imaging. 2023;16(9):1181-1189.

**EFFECTIVE DATE** 7/31/2026

ID\_CHS\_2026