

Medical Policy Manual **Approved Rev: Do Not Implement until 9/30/26**

Epcoritamab-bysp (Epkinly™)

IMPORTANT REMINDER

We develop Medical Policies to provide guidance to Members and Providers. This Medical Policy relates only to the services or supplies described in it. The existence of a Medical Policy is not an authorization, certification, explanation of benefits or a contract for the service (or supply) that is referenced in the Medical Policy. For a determination of the benefits that a Member is entitled to receive under his or her health plan, the Member's health plan must be reviewed. If there is a conflict between the medical policy and a health plan or government program (e.g., TennCare), the express terms of the health plan or government program will govern.

POLICY

INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

- Epkinly is indicated for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma (DLBCL) not otherwise specified, including DLBCL arising from indolent lymphoma, and high-grade B-cell lymphoma (HGBL), after two or more lines of systemic therapy.
- **Epkinly is indicated for the treatment of adult patients with relapsed or refractory follicular lymphoma (FL) in combination with lenalidomide and rituximab.**
- Epkinly is indicated for the treatment of adult patients with relapsed or refractory follicular lymphoma (FL) **as monotherapy** after two or more lines of systemic therapy.

Compendial Uses

- Diffuse Large B-Cell Lymphomas
- High Grade B-Cell Lymphomas
- Histologic Transformation of Indolent Lymphomas to Diffuse Large B-Cell Lymphoma
- Human Immunodeficiency Virus (HIV)- Related B-Cell Lymphomas
- Monomorphic Post-Transplant Lymphoproliferative Disorders
- Follicular Lymphoma
- **Histologic (Richter) transformation**

All other indications are considered experimental/investigational and not medically necessary.

COVERAGE CRITERIA

B-Cell Lymphomas

Authorization of 12 months may be granted for treatment of B-cell lymphoma when **one** of the following criteria is met:

- **The requested medication will be used as subsequent therapy for follicular lymphoma in combination with lenalidomide and rituximab for up to 12 cycles total for no response, relapsed, or progressive disease.**
- The requested medication will be used as subsequent therapy in combination with GemOx (gemcitabine and oxaliplatin) for relapsed/refractory disease with any of the following subtypes:



Medical Policy Manual **Approved Rev: Do Not Implement until 9/30/26**

- Diffuse large B-cell lymphoma (DLBCL)
- High grade B- cell lymphoma
- HIV-related B- cell lymphoma including HIV-related DLBCL, primary effusion lymphoma, HHV8-positive DLBCL, not otherwise specified, and HIV-related plasmablastic lymphoma
- Monomorphic post-transplant lymphoproliferative disorder (B-cell type)
- The requested medication will be used as a single agent after at least 2 prior lines of systemic therapy when the member has partial response, no response, progressive, relapsed or refractory disease with any of the following subtypes:
 - Diffuse large B-cell lymphoma (DLBCL)
 - High grade B-cell lymphoma
 - Histologic transformation of indolent lymphoma to DLBCL
 - HIV-related B-cell lymphoma including HIV-related DLBCL, primary effusion lymphoma, and HHV8-positive DLBCL, not otherwise specified
 - Monomorphic post-transplant lymphoproliferative disorder (B-cell type)
 - Follicular lymphoma

Histologic (Richter) Transformation to DLBCL

Authorization of 12 months may be granted for treatment of histologic (Richter) transformation to DLBCL.

CONTINUATION OF THERAPY

Authorization of 12 months may be granted for continued treatment in members requesting reauthorization for an indication listed in the coverage criteria section when there is no evidence of unacceptable toxicity or disease progression while on the current regimen. **A maximum of 12 cycles total will be authorized for members using the requested medication in combination with lenalidomide and rituximab for follicular lymphoma.**

APPLICABLE TENNESSEE STATE MANDATE REQUIREMENTS


BlueCross BlueShield of Tennessee's Medical Policy complies with Tennessee Code Annotated Section 56-7-2352 regarding coverage of off-label indications of Food and Drug Administration (FDA) approved drugs when the off-label use is recognized in one of the statutorily recognized standard reference compendia or in the published peer-reviewed medical literature.

ADDITIONAL INFORMATION

For appropriate chemotherapy regimens, dosage information, contraindications, precautions, warnings, and monitoring information, please refer to one of the standard reference compendia (e.g., the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) published by the National Comprehensive Cancer Network®, Drugdex Evaluations of Micromedex Solutions at Truven Health, or The American Hospital Formulary Service Drug Information).

REFERENCES

1. Epkinly [package insert]. Plainsboro, NJ: Genmab US, Inc.; **March 2026**
2. The NCCN Drugs & Biologics Compendium® © 2026 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed April 9, 2026.



Policy

Medical Policy Manual **Approved Rev: Do Not Implement until 9/30/26**

EFFECTIVE DATE 9/30/2026

ID_CHS_2025a