



Medical Policy Manual **Approved Rev: Do Not Implement until 12/1/26**

Leuprolide Suspension (Lupron Depot®, Leuprolide Acetate Depot 1-Month 3.75mg, 3-Month 11.25 mg)

IMPORTANT REMINDER

We develop Medical Policies to provide guidance to Members and Providers. This Medical Policy relates only to the services or supplies described in it. The existence of a Medical Policy is not an authorization, certification, explanation of benefits or a contract for the service (or supply) that is referenced in the Medical Policy. For a determination of the benefits that a Member is entitled to receive under his or her health plan, the Member's health plan must be reviewed. If there is a conflict between the medical policy and a health plan or government program (e.g., TennCare), the express terms of the health plan or government program will govern.

POLICY

INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

FDA-Approved Indications

Endometriosis

Lupron Depot 3.75 mg and Lupron Depot 3-Month 11.25 mg are indicated for management of endometriosis, including pain relief and reduction of endometriotic lesions. Lupron Depot 3.75 mg and Lupron Depot 3-Month 11.25 mg **in combination** with norethindrone acetate 5 mg daily are also indicated for initial management of the painful symptoms of endometriosis and for management of recurrence of symptoms.

Use of norethindrone acetate in combination with Lupron Depot 3.75 mg and Lupron Depot **3-Month** 11.25 mg is referred to as add-back therapy, and is intended to reduce the loss of bone mineral density (BMD) and reduce vasomotor symptoms associated with use of Lupron Depot 3.75 mg and Lupron Depot **3-Month** 11.25 mg.

Uterine Leiomyomata (Fibroids)

When used concomitantly with iron therapy, Lupron Depot 3.75 mg and Lupron Depot-3 Month 11.25 mg are indicated for preoperative hematologic improvement of women with anemia caused by fibroids for whom three months of hormonal suppression is deemed necessary. Consider a one-month trial period on iron alone, as some women will respond to iron alone. Lupron Depot may be added if the response to iron alone is considered inadequate.

Limitations of Use

For endometriosis: The total duration of therapy with Lupron Depot 3.75 mg and Lupron Depot 3-Month 11.25 mg plus add-back therapy should not exceed 12 months due to concerns about adverse impact on bone mineral density.

For uterine leiomyomata (fibroids): Lupron Depot 3.75 mg and **Lupron Depot 3-Month** 11.25 mg are not indicated for combination use with norethindrone acetate add-back therapy for the preoperative hematologic improvement of women with anemia caused by heavy menstrual bleeding due to fibroids.

Compendial Uses

- Breast cancer

Medical Policy Manual **Approved Rev: Do Not Implement until 12/1/26**

- Ovarian cancer
 - Epithelial ovarian cancer/fallopian tube cancer/primary peritoneal cancer
 - Carcinosarcoma (malignant mixed Müllerian tumors)
 - Grade 1 endometrioid carcinoma
 - Low-grade serous carcinoma
 - Mucinous carcinoma of the ovary
 - Clear cell carcinoma of the ovary
- Salivary gland tumor
- Gender dysphoria (also known as transgender and gender diverse [TGD] persons)
- Preservation of ovarian function
- Prevention of recurrent menstrual related attacks in acute porphyria
- Uterine Sarcoma

All other indications are considered experimental/investigational and not medically necessary.

DOCUMENTATION

Submission of the following information is necessary to initiate the prior authorization review, where applicable, test results confirming:

- Androgen receptor (AR) positive status
- Hormone receptor (HR) positive status

PRESCRIBER SPECIALTIES

Gender Dysphoria

The medication must be prescribed by or in consultation with a provider specialized in the care of transgender youth (e.g., pediatric endocrinologist, family or internal medicine physician, obstetrician-gynecologist) that has collaborated care with a mental health provider for members less than 18 years of age.

Prevention of Recurrent Menstrual Related Attacks in Acute Porphyria

The medication must be prescribed by or in consultation with a provider experienced in the management of porphyrias.

COVERAGE CRITERIA

Endometriosis

Authorization of up to 6 months (one treatment course) may be granted to members for initial treatment of endometriosis.

Uterine Leiomyomata (Fibroids)

Authorization of up to 3 months may be granted for initial treatment of uterine leiomyomata (fibroids) when either of the following criteria is met:

- Member has anemia due to uterine leiomyomata
- Lupron Depot will be used prior to surgery for uterine leiomyomata.

Breast Cancer

Medical Policy Manual **Approved Rev: Do Not Implement until 12/1/26**

Authorization of 12 months may be granted for treatment of hormone receptor-positive breast cancer.

Ovarian Cancer

Authorization of 12 months may be granted for treatment of persistent disease or recurrence of any of the following types of ovarian cancer when used as a single agent:

- Epithelial ovarian cancer
- Fallopian tube cancer
- Primary peritoneal cancer
- Grade 1 endometrioid carcinoma
- Low-grade serous carcinoma
- Carcinosarcoma (malignant mixed Müllerian tumors)
- Mucinous carcinoma of the ovary
- Clear cell carcinoma of the ovary

Salivary Gland Tumors

Authorization of 12 months may be granted for treatment of recurrent, unresectable, or metastatic salivary gland tumor as a single agent or in combination with abiraterone and prednisone when the tumor is androgen receptor positive.

Gender Dysphoria

*Individual is age 18 or older or the individual is less than age 18 as permissive under applicable law

Authorization of 12 months may be granted for pubertal hormonal suppression in an adolescent member **who is seeking gender-affirming care** when all of the following criteria are met:

- The member has a diagnosis of gender dysphoria.
- The member is able to make an informed decision to engage in treatment.
- The member has reached Tanner stage 2 of puberty or greater.
- The member's comorbid conditions are reasonably controlled.
- The member has been educated on any contraindications and side effects to therapy.
- The member has been informed of fertility preservation options.

Authorization of 12 months may be granted for gender transition **in a member who is seeking gender-affirming care** when all of the following criteria are met:

- The member has a diagnosis of gender dysphoria.
- The member is able to make an informed decision to engage in treatment.
- The member will receive the requested medication concomitantly with gender-affirming hormones.
- The member's comorbid conditions are reasonably controlled.
- The member has been educated on any contraindications and side effects to therapy.
- The member has been informed of fertility preservation options.

Preservation of Ovarian Function

Authorization of 3 months may be granted for preservation of ovarian function when the member is premenopausal and undergoing chemotherapy.

Prevention of Recurrent Menstrual Related Attacks in Acute Porphyria

Authorization of 12 months may be granted for prevention of recurrent menstrual related attacks in members with acute porphyria.

Uterine Sarcoma



Medical Policy Manual **Approved Rev: Do Not Implement until 12/1/26**

Authorization of 12 months may be granted for treatment of uterine sarcoma in combination with an aromatase inhibitor (e.g., anastrozole, exemestane, letrozole) when the member is premenopausal and not suitable for surgery.

CONTINUATION OF THERAPY

Endometriosis

Authorization of up to 6 months (for a lifetime maximum of 12 months total) may be granted for retreatment of endometriosis when both of the following criteria are met:

- The member has had a recurrence of symptoms.
- The member has a bone mineral density within normal limits.

Uterine Leiomyomata (fibroids)

Authorization of up to 3 months (for a lifetime maximum of 6 months total) may be granted when either of the following criteria is met:

- Member has anemia due to uterine leiomyomata
- Lupron Depot will be used prior to surgery for uterine leiomyomata.

Breast Cancer, Ovarian Cancer, Salivary Gland Tumor, and Uterine Sarcoma

Authorization of 12 months may be granted for continued treatment of breast cancer, ovarian cancer, salivary gland tumor, and uterine sarcoma in members requesting reauthorization when there is no evidence of unacceptable toxicity or disease progression while on the current regimen.

Gender Dysphoria

*Individual is age 18 or older or the individual is less than age 18 as permissive under applicable law

Authorization of 12 months may be granted for continued treatment for pubertal hormonal suppression in an adolescent member **who is seeking gender-affirming care and** requesting reauthorization when all of the following criteria are met:

- The member has a diagnosis of gender dysphoria.
- The member is able to make an informed decision to engage in treatment.
- The member has previously reached Tanner stage 2 of puberty or greater.
- The member's comorbid conditions are reasonably controlled.
- The member has been educated on any contraindications and side effects to therapy.
- Before the start of therapy, the member has been informed of fertility preservation options.

Authorization of 12 months may be granted for continued treatment for gender transition in a member **who is seeking gender-affirming care and** requesting reauthorization when all of the following criteria are met:

- The member has a diagnosis of gender dysphoria.
- The member is able to make an informed decision to engage in treatment.
- The member will receive requested medication concomitantly with gender-affirming hormones.
- The member's comorbid conditions are reasonably controlled.
- The member has been educated on any contraindications and side effects to therapy.
- Before the start of therapy, the member has been informed of fertility preservation options.

Preservation of Ovarian Function and Prevention of Recurrent Menstrual Related Attacks in Acute Porphyria

Medical Policy Manual **Approved Rev: Do Not Implement until 12/1/26**

All members (including new members) requesting authorization for continuation of therapy must meet all the requirements in the coverage criteria section.

OTHER

Per state regulatory guidelines around gender dysphoria, age restrictions may apply.

APPLICABLE TENNESSEE STATE MANDATE REQUIREMENTS

BlueCross BlueShield of Tennessee's Medical Policy complies with Tennessee Code Annotated Section 56-7-2352 regarding coverage of off-label indications of Food and Drug Administration (FDA) approved drugs when the off-label use is recognized in one of the statutorily recognized standard reference compendia or in the published peer-reviewed medical literature.

ADDITIONAL INFORMATION

For appropriate chemotherapy regimens, dosage information, contraindications, precautions, warnings, and monitoring information, please refer to one of the standard reference compendia (e.g., the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) published by the National Comprehensive Cancer Network®, Drugdex Evaluations of Micromedex Solutions at Truven Health, or The American Hospital Formulary Service Drug Information).

REFERENCES

1. Lupron Depot 3.75 mg [package insert]. North Chicago, IL: AbbVie Inc.; **September 2025**.
2. Lupron Depot-3 Month 11.25 mg [package insert]. North Chicago, IL: AbbVie Inc.; **September 2025**.
3. The NCCN Drugs & Biologics Compendium® © 2026 National Comprehensive Cancer Network, Inc. <http://www.nccn.org>. Accessed **February 10, 2026**.
4. Management of symptomatic uterine leiomyomas: ACOG Practice Bulletin No. 228. American College of Obstetricians and Gynecologists. *Obstet Gynecol*. 2021 June 1;137(6):e100-e115.
5. Marret H, Fritel X, Ouldamer L, et al. Therapeutic management of uterine fibroid tumors: updated French guidelines. *European Journal of Obstetrics and Gynecology and Reproductive Biology*. 2012;165:156-164.
6. Hembree WC, Cohen-Kettenis PT, Gooren L, et al. Endocrine treatment of gender-dysphoric/gender-incongruent persons: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2017;102(11):3869-3903.
7. Coleman E, Radix AE, Brown GR, et al. Standards of care for the health of transgender and gender diverse people, version 8. 2022;23 (Suppl 1): S1-S259. doi: 10.1080/26895269.2022.2100644
8. Oktay K, Harvey BE, Patridge AH, et al. Fertility preservation in patients with cancer: ASCO Clinical Practice Guideline Update. *J Clin Oncol*. 2018 Jul 1;36(19):1994-2001.
9. Clowse MEB, Behera MA, Anders CK, et al. Ovarian preservation by GnRH agonists during chemotherapy: a meta-analysis. *J Womens Health (Larchmt)*. 2009 Mar; 18(3): 311–319. doi:10.1089/jwh.2008.0857
10. Stein P, Badminton M, Barth J, et al. British and Irish Porphyria Network. Best practice guidelines on clinical management of acute attacks of porphyria and their complications. *Ann Clin Biochem*. 2013 May;50(Pt 3):217-23.
11. Innala E, Bäckström T, Bixo M, et al. C. Evaluation of gonadotrophin-releasing hormone agonist treatment for prevention of menstrual-related attacks in acute porphyria. *Acta Obstet Gynecol* 2010;89:95–100.
12. Mahfouda S, Moore JK, Siafarikas A, et al. Puberty suppression in transgender children and adolescents. *Lancet Diabetes Endocrinol*. 2017; 5(10): 816-826.
13. **American College of Obstetricians and Gynecologists' Committee on Gynecologic Practice; American College of Obstetricians and Gynecologists' Committee on Health Care for Underserved Women. Health care for**

Medical Policy Manual **Approved Rev: Do Not Implement until 12/1/26**

transgender and gender diverse individuals: ACOG Committee opinion, number 823. Obstet Gynecol. 2021;137(3):e75-e88.

14. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Head and Neck Cancers. Version 1.2026. Accessed February 13, 2026. https://www.nccn.org/professionals/physician_gls/pdf/head-and-neck.pdf.

EFFECTIVE DATE 12/1/2026

ID_CHS_60492026_0926